### THIS TEST BOOKLET MAY NOT BE TAKEN FROM THE TESTING ROOM D&SDT MOCK TEST TEMPLATE Transcription Test #777

### General Instructions

You have taken Mack Testmann to the health care provider and have received the medication from the pharmacy. The date is November 15, yr. The time is 1:00pm.

Use the health care provider's order, pharmacy label and generic equivalents to discontinue the appropriate medication order and transcribe the new medication order onto the medication sheet.

You are not being asked to administer this medication. This is a transcription exercise only.

Only the marks you make on your medication sheet will be scored.

When you have finished, raise your hand and the test administrator will collect the test and all testing materials.

# HEALTH CARE PROVIDER ORDER D&SDT MOCK TEST TEMPLATE Transcription Test #777

Date: November 15, yr		
Allergies: Seafood		
Reason for visit: Mack Testmann began swimming lessons is rubbing left ear.		
Enteric 160 mg by mouth twice a day.		
Glidden 75 mg by mouth once a day in morning.		
Collate 180 mg by mouth twice a day.		
Date: November 15, yr		
Health Care Provider Findings:		
Left ear infection		
New / Changed Medication / Treatment Orders:		
Discontinue Enteric 160 mg by mouth twice a day.		
Begin Immunal 250 mg by mouth twice a day for ten days.		
Instructions:		
Lab Work or Tests: None		
Date: November 15, yr		

## PHARMACY LABEL D&SDT MOCK TEST TEMPLATE Transcription Test #777

Rx # 777-7722 Pills Pharmacy
1214 Main Street
Waterfield, MA 02422

Mack Testmann November 15, yr.

Immunal 250 mg
IC: Toraminal 250 mg
Take one tablet by mouth twice a day for ten days.

Dr. Mills

Lot # 4132-56 Exp Date: 11-15-yr. Refills: 0

### **GENERIC EQUIVALENTS**

Brand	Generic
Collate	Cusate Clorate
Enteric	Hydroxzine Hydrosodium
Immunal	Toraminal
Glidden	Motrigine
Artanal	Trihexyphenidyl Hydroclorate

### **MEDICATION INFORMATION**

Immunal is an antibiotic used to treat inner ear infections.

Common side effects include lethargy, coughing, and loose stools. Finish all medication. Take with food.